



# OFF-MOMENTUM PARTICLES

---

Eric Prebys, UC Davis



USPAS Fundamentals, June 4-15, 2018 E. Prebys, Accelerator Fundamentals: Off-Momentum Particles 2 

## Off-Momentum Particles

- Our previous discussion implicitly assumed that all particles were at the same momentum
  - Each quad has a constant focal length
  - There is a single nominal trajectory
- In practice, this is never true. Particles will have a distribution about the nominal momentum
- We will characterize the behavior of off-momentum particles in the following ways
  - “Dispersion” ( $D$ ): the dependence of position on deviations from the nominal momentum
 
$$\Delta x(s) = D_x(s) \frac{\Delta p}{p_0}$$

$D$  has units of length
  - “Chromaticity” ( $\eta$ ): the change in the tune caused by the different focal lengths for off-momentum particles
 
$$\Delta \nu_x = \xi_x \frac{\Delta p}{p_0} \left( \text{sometimes } \frac{\Delta \nu_x}{\nu_x} = \xi_x \frac{\Delta p}{p_0} \right)$$
  - Path length changes (momentum compaction)
 
$$\frac{\Delta L}{L} = \alpha \frac{\Delta p}{p}$$


USPAS Fundamentals, June 4-15, 2018 E. Prebys, Accelerator Fundamentals: Off-Momentum Particles 3

## Review: Equations of Motion

- Recall that in a curvilinear coordinate system, the equations of motion become

$$x'' = -\frac{B_y}{(B\rho)} \left(1 + \frac{x}{\rho}\right)^2 + \frac{\rho+x}{\rho^2}$$

$$y'' = \frac{B_x}{(B\rho)} \left(1 + \frac{x}{\rho}\right)^2$$



Note:  $s$  measured along nominal trajectory,  $v_s$  measured along actual trajectory

$$\dot{s} = \frac{ds}{dt} = \frac{\rho}{r} v_s$$

- We'll now consider the effect of off momentum particle by comparing the "true" rigidity to the nominal rigidity

$$(B\rho)_{true} = (B\rho) \frac{p}{p_0} \rightarrow \frac{1}{(B\rho)_{true}} = \frac{1}{(B\rho)} \frac{p_0}{p} = \frac{1}{(B\rho)} \frac{p_0}{p_0 + \Delta p} \approx \frac{1}{(B\rho)} \left(1 - \frac{\Delta p}{p_0}\right)$$

USPAS Fundamentals, June 4-15, 2018 E. Prebys, Accelerator Fundamentals: Off-Momentum Particles 4

## Off-Momentum Particles

- If we substitute this into the equations of motion, and keep only linear terms, we end up with one new term in each equation

$$x'' = -\frac{B_y}{(B\rho)} \left(1 - \frac{\Delta p}{p_0}\right) \left(1 + \frac{x}{\rho}\right)^2 + \frac{\rho+x}{\rho^2} = (\dots) + \frac{B_y}{(B\rho)} \frac{\Delta p}{p_0} \equiv (\dots) + \frac{B_y}{(B\rho)} \delta$$

$$y'' = \frac{B_x}{(B\rho)} \left(1 - \frac{\Delta p}{p_0}\right) \left(1 + \frac{x}{\rho}\right)^2 = (\dots) - \frac{B_x}{(B\rho)} \delta$$

- The parts in parentheses just give us our nominal equations of motion. We now invoke
 
$$B_x = B' y \approx 0$$

$$B_y = B_0 + B' x \approx B_0;$$

$$\frac{B_y}{(B\rho)} \approx \frac{B_0}{(B\rho)} = \frac{1}{\rho}$$

$$\frac{B_x}{(B\rho)} \approx 0$$

- And our new equations become

$$x'' + \left( \frac{1}{\rho^2} + \frac{1}{(B\rho)} B' \right) x = \frac{1}{\rho} \delta; \quad y'' - \frac{1}{(B\rho)} B' y = 0$$

New

USPAS Fundamentals, June 4-15, 2018 E. Prebys, Accelerator Fundamentals: Off-Momentum Particles 5

- This is a second order differential inhomogeneous differential equation, so the solution is
 
$$x(s) = x_0 C(s) + x'_0 S(s) + \delta d(s)$$

$$x'(s) = x_0 C'(s) + x'_0 S'(s) + \delta d'(s)$$

Where  $d(s)$  is the solution particular solution of the differential equation

$$d'' + Kd = \frac{1}{\rho}$$

- We solve this piecewise, for  $K$  constant and find
 
$$K > 0: \quad d(s) = \frac{1}{\rho K} (1 - \cos \sqrt{K} s)$$

$$d'(s) = \frac{1}{\rho \sqrt{K}} \sin \sqrt{K} s$$

$$K < 0: \quad d(s) = -\frac{1}{\rho K} (1 - \cosh \sqrt{K} s)$$

$$d'(s) = \frac{1}{\rho \sqrt{K}} \sinh \sqrt{K} s$$

USPAS Fundamentals, June 4-15, 2018 E. Prebys, Accelerator Fundamentals: Off-Momentum Particles 6

## General Solution

- The general solution is now
 
$$x(s) = x_0 C(s) + x'_0 S(s) + \delta d(s)$$

$$x'(s) = x_0 C'(s) + x'_0 S'(s) + \delta d'(s)$$

Solution to the on-momentum case
Off-momentum correction
- We can express this in matrix form as
 

Usual transfer matrix

$$\begin{pmatrix} x(s) \\ x'(s) \\ \delta \end{pmatrix} = \begin{pmatrix} m_{11} & m_{12} \\ m_{21} & m_{22} \\ 0 & 0 \end{pmatrix} \begin{pmatrix} d(s) \\ d'(s) \\ 1 \end{pmatrix} \begin{pmatrix} x_0 \\ x'_0 \\ \delta \end{pmatrix}$$

USPAS Fundamentals, June 4-15, 2018 E. Prebys, Accelerator Fundamentals: Off-Momentum Particles 7

## New Equilibrium Orbit

- We want to solve for an orbit of an off-momentum particle that follows the periodicity of the machine.
- This will serve as the new equilibrium orbit for off-momentum particles.

$x(s, \delta) = D_x(s) \delta$ 
“Dispersion” [L]

- This must satisfy

$$\begin{pmatrix} \delta D_x \\ \delta D'_x \\ \delta \end{pmatrix} = \begin{pmatrix} M_{11} & M_{12} & d \\ M_{21} & M_{22} & d' \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} \delta D_x \\ \delta D'_x \\ \delta \end{pmatrix} \Rightarrow \begin{pmatrix} D_x \\ D'_x \\ 1 \end{pmatrix} = \begin{pmatrix} \dots \\ D'_x \\ 1 \end{pmatrix}$$

USPAS Fundamentals, June 4-15, 2018 E. Prebys, Accelerator Fundamentals: Off-Momentum Particles 8

## Simplifying Assumptions

- For the most part, we will consider systems for which both of the following are true
  - “separated function”: Separate dipoles and quadrupoles
 
$$\Rightarrow \frac{1}{\rho^2} \text{ and } B' \text{ are never both non-zero at the same point}$$
  - “Isomagnetic”: All bend dipoles have the same field
 
$$\frac{1}{\rho^2} = \frac{1}{\rho_0^2} \text{ inside of bend dipoles}$$

$$= 0 \text{ everywhere else}$$

USPAS Fundamentals, June 4-15, 2018 E. Prebys, Accelerator Fundamentals: Off-Momentum Particles 9

### Example: FODO Cell

- We look at our symmetric FODO cell, but assume that the drifts are bend magnets that take up the entire space (a pretty good assumption)
- Each bends the beam by an angle  $\theta$

For a thin lens  $d \sim d' \sim 0$ . For a pure bend magnet  $s \ll \rho_0$

$$K = \frac{1}{\rho_0^2}: \quad d(L) = \frac{1}{\rho_0 K} (1 - \cos \sqrt{KL}) = \rho_0 \left( 1 - \cos \frac{L}{\rho_0} \right) \approx \frac{1}{2\rho_0} L^2 \rightarrow \frac{1}{2} \frac{L^2}{\rho_0} = \frac{1}{2} \theta L$$

$$d'(L) = \frac{1}{\rho_0 \sqrt{K}} \sin \sqrt{KL} = \sin \frac{L}{\rho_0} \approx \frac{L}{\rho_0} \rightarrow \theta$$

USPAS Fundamentals, June 4-15, 2018 E. Prebys, Accelerator Fundamentals: Off-Momentum Particles 10

### Transfer Matrix

- We put this all together to get a transfer matrix of the form

$$\begin{pmatrix} x(s) \\ x'(s) \\ \delta \end{pmatrix} = \begin{pmatrix} M_{11} & M_{12} \\ M_{21} & M_{22} \\ 0 & 0 \end{pmatrix} \begin{pmatrix} d(s) \\ d'(s) \\ 1 \end{pmatrix} \begin{pmatrix} x_0 \\ x'_0 \\ \delta \end{pmatrix}$$

Usual transfer matrix

- Using our solutions from the previous page, we get

$$M = \begin{pmatrix} 1 & 0 & 0 \\ -\frac{1}{2f} & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & L & \frac{L\theta}{2} \\ 0 & 1 & \theta \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ \frac{1}{f} & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & L & \frac{L\theta}{2} \\ 0 & 1 & \theta \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ -\frac{1}{2f} & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

- For a ring:

$$\theta = \frac{2\pi}{2N_{cell}} = \frac{\pi}{N_{cell}} = \begin{pmatrix} 1 - \frac{L^2}{2f^2} & 2L \left( 1 + \frac{L}{2f} \right) & 2L\theta \left( 1 + \frac{L}{4f} \right) \\ -\frac{L}{2f^2} + \frac{L^2}{4f^3} & 1 - \frac{L^2}{2f^2} & 2\theta \left( 1 - \frac{L}{4f} - \frac{L^2}{8f^2} \right) \\ 0 & 0 & 1 \end{pmatrix}$$

USPAS Fundamentals, June 4-15, 2018 E. Prebys, Accelerator Fundamentals: Off-Momentum Particles 11

## Solving for Dispersion

- We must solve

$$\begin{pmatrix} D \\ D' \\ 1 \end{pmatrix} = \begin{pmatrix} 1 - \frac{L^2}{2f^2} & 2L\left(1 + \frac{L}{2f}\right) & 2L\theta\left(1 + \frac{L}{4f}\right) \\ -\frac{L}{2f^2} + \frac{L^2}{4f^3} & 1 - \frac{L^2}{2f^2} & 2\theta\left(1 - \frac{L}{4f} - \frac{L^2}{8f^2}\right) \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} D \\ D' \\ 1 \end{pmatrix}$$

- In your homework, you show that

$$D_{F,D} = \frac{\theta L \left(1 \pm \frac{1}{2} \sin \frac{\mu}{2}\right)}{\sin^2 \frac{\mu}{2}}$$

USPAS Fundamentals, June 4-15, 2018 E. Prebys, Accelerator Fundamentals: Off-Momentum Particles 12

## Evolution of Dispersion Functions

- Since the dispersion functions represent displacements, they will evolve like the position

$$\begin{pmatrix} D_x(s) \\ D'_x(s) \\ 1 \end{pmatrix} = \begin{pmatrix} m_{11} & m_{12} & d(s) \\ m_{21} & m_{22} & d'(s) \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} D_x(0) \\ D'_x(0) \\ 1 \end{pmatrix}$$

- Putting it all together

USPAS Fundamentals, June 4-15, 2018 E. Prebys, Accelerator Fundamentals: Off-Momentum Particles 13

## Momentum Compaction Factor

- In general, particles with a high momentum will travel a longer path length. We have

$$C(p_0) = \oint ds$$

$$C(p_0 + \Delta p) = \oint \left( 1 + \frac{D \Delta p}{\rho p_0} \right) ds$$

$$\frac{\Delta C}{C} = \frac{\oint \frac{D}{\rho} ds}{\oint ds} \delta = \left\langle \frac{D}{\rho} \right\rangle \delta$$

"momentum compaction" factor  $\Rightarrow \alpha_c \delta$

So yes, we now have an ambiguous definition of  $\alpha$ , too!

USPAS Fundamentals, June 4-15, 2018 E. Prebys, Accelerator Fundamentals: Off-Momentum Particles 14

## Slip Factor

- The "slip factor" is defined as the fractional change in the orbital period divided by the fractional change in momentum

$$T = \frac{C}{v}$$

$$\frac{\Delta T}{T} = \frac{\Delta C}{C} - \frac{\Delta v}{v} = \frac{\Delta C}{C} - \frac{\Delta \beta}{\beta}$$

$$= \alpha \frac{\Delta p}{p} - \frac{1}{\gamma^2} \frac{\Delta p}{p}$$

$$= \left( \alpha - \frac{1}{\gamma^2} \right) \frac{\Delta p}{p}$$

$$\equiv \eta \frac{\Delta p}{p}$$

$\gamma < \frac{1}{\sqrt{\alpha}}: \eta < 0$  velocity dominates

$\gamma > \frac{1}{\sqrt{\alpha}}: \eta > 0$  momentum dominates

$\gamma = \frac{1}{\sqrt{\alpha}}: \eta = 0$  "transition"

Transition gamma or "gamma-T"  $\rightarrow \gamma_T \equiv \frac{1}{\sqrt{\alpha}} \Rightarrow \eta = \left( \frac{1}{\gamma_T^2} - \frac{1}{\gamma^2} \right)$

USPAS Fundamentals, June 4-15, 2018 E. Prebys, Accelerator Fundamentals: Off-Momentum Particles 15

## Special Cases for Slip Factor

- Linacs:  $\alpha=0 \rightarrow \eta = -\frac{1}{\gamma^2}$  (always negative)
- Simple Cyclotrons:  $C = 2\pi\rho = 2\pi \frac{p}{eB} \rightarrow \alpha=1 \rightarrow \eta = \left(1 - \frac{1}{\gamma^2}\right)$  (0 to positive)
- Synchrotrons: more complicated
  - Negative below  $\gamma_T$
  - Positive above  $\gamma_T$

$$\eta = \left(\alpha - \frac{1}{\gamma^2}\right)$$

USPAS Fundamentals, June 4-15, 2018 E. Prebys, Accelerator Fundamentals: Off-Momentum Particles 16

## Transition $\gamma$ for Synchrotrons (approx.)

- For a simple FODO CELL
 
$$\beta_{\max, \min} = 2L \frac{\left(1 \pm \sin \frac{\mu}{2}\right)}{\sin \mu}; \text{ and } D_{\max, \min} = \theta L \frac{\left(1 \pm \frac{1}{2} \sin \frac{\mu}{2}\right)}{\sin^2 \frac{\mu}{2}}$$
- If we assume they vary ~linearly between maxima, then for small  $\mu$ 

$$\langle \beta \rangle \approx \frac{2L}{\mu}; \quad \langle D \rangle \approx \frac{4\theta L}{\mu^2} = 4 \frac{L^2}{\mu^2 \rho} = \frac{\langle \beta \rangle^2}{\rho}$$
- Also
 
$$v = \frac{1}{2\pi} \oint \frac{ds}{\beta(s)} \approx \frac{1}{2\pi} \frac{2\pi R}{\langle \beta \rangle} \approx \frac{\rho}{\langle \beta \rangle}$$



USPAS Fundamentals, June 4-15, 2018 E. Prebys, Accelerator Fundamentals: Off-Momentum Particles 17

(cont'd)

- We just showed  $\langle D \rangle \approx \frac{\langle \beta \rangle^2}{\rho}$   $\rightarrow$   $\langle D \rangle \approx \frac{\rho}{v^2}$   
 $v \approx \frac{\rho}{\langle \beta \rangle}$
- So  $\alpha_c = \frac{1}{C} \oint \frac{D}{\rho} ds \approx \frac{1}{\rho} \langle D \rangle \approx \frac{1}{v^2}$   
 $\gamma_t = \frac{1}{\sqrt{\alpha_c}} \approx v$
- This approximation generally works better than it should
  - FNAL Booster:  $v=6.8, \gamma_T=5.5$

USPAS Fundamentals, June 4-15, 2018 E. Prebys, Accelerator Fundamentals: Off-Momentum Particles 18

### Digression: Quadrupole Perturbation

- We can express the matrix for a complete revolution at a point as
 
$$\mathbf{M}(s) = \begin{pmatrix} \cos 2\pi\nu + \alpha(s) \sin 2\pi\nu & \beta(s) \sin 2\pi\nu \\ -\gamma(s) \sin 2\pi\nu & \cos 2\pi\nu - \alpha(s) \sin 2\pi\nu \end{pmatrix}$$
- If we add focusing quad at this point, we have
 
$$\mathbf{M}'(s) = \begin{pmatrix} \cos 2\pi\nu_0 + \alpha(s) \sin 2\pi\nu_0 & \beta(s) \sin 2\pi\nu_0 \\ -\gamma(s) \sin 2\pi\nu_0 & \cos 2\pi\nu_0 - \alpha(s) \sin 2\pi\nu_0 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ -\frac{1}{f} & 1 \end{pmatrix}$$

$$= \begin{pmatrix} \cos 2\pi\nu_0 + \alpha(s) \sin 2\pi\nu_0 - \frac{\beta(s)}{f} \sin 2\pi\nu_0 & \beta(s) \sin 2\pi\nu_0 \\ -\gamma(s) \sin 2\pi\nu_0 - \frac{1}{f} (\cos 2\pi\nu_0 - \alpha(s) \sin 2\pi\nu_0) & \cos 2\pi\nu_0 - \alpha(s) \sin 2\pi\nu_0 \end{pmatrix}$$
- We calculate the trace to find the new tune
 
$$\cos 2\pi\nu = \frac{1}{2} \text{Tr}(\mathbf{M}') = \cos 2\pi\nu_0 - \frac{1}{2f} \beta(s) \sin 2\pi\nu_0$$
- For small changes  $\cos 2\pi(\nu_0 + \Delta\nu) \approx \cos 2\pi\nu_0 - 2\pi \sin 2\pi\nu_0 \Delta\nu = \cos 2\pi\nu_0 - \frac{1}{2f} \beta(s) \sin 2\pi\nu_0$   
 $\Rightarrow \Delta\nu = \frac{1}{4\pi} \frac{\beta(s)}{f}$

USPAS Fundamentals, June 4-15, 2018 E. Prebys, Accelerator Fundamentals: Off-Momentum Particles 19

## Total Tune Shift

- The focal length associated with a local anomalous gradient is

$$d\left(\frac{1}{f}\right) = \frac{B'}{(B\rho)} ds$$

- So the total tune shift is

$$\Delta\nu = \frac{1}{4\pi} \oint \beta(s) \frac{B'(s)}{(B\rho)} ds$$

USPAS Fundamentals, June 4-15, 2018 E. Prebys, Accelerator Fundamentals: Off-Momentum Particles 20

## Chromaticity

- In general, momentum changes will lead to a tune shift by changing the effective focal lengths of the magnets
- We already showed

$$\frac{1}{f} = \frac{B'l}{(B\rho)} = \frac{B'l}{(B\rho)_0} \frac{p_0}{p} \approx \frac{1}{f_0} \left(1 - \frac{\Delta p}{p_0}\right)$$

$$\Rightarrow \Delta\nu = -\frac{1}{4\pi} \sum_i \beta_i \frac{1}{f_i} \frac{\Delta p}{p_0} \equiv \xi \frac{\Delta p}{p_0}$$

- Where

$$\frac{1}{f_0} = -\int_0^L \frac{B'}{(B\rho)} ds$$

USPAS Fundamentals, June 4-15, 2018 E. Prebys, Accelerator Fundamentals: Off-Momentum Particles 21

## Chromaticity (Cont'd)

- Recalling that in our general equation of motion
 
$$x'' + \left( \frac{1}{\rho^2} + \frac{B'(s)}{B\rho} \right) x = 0 \equiv x'' + K(s)x$$
- We see that the effective focal length for a region is
 
$$\frac{1}{f_0} = \int_0^L \frac{B'}{B\rho} ds \Rightarrow \frac{1}{f_{eff}} = \int_0^L \left( \frac{1}{\rho^2} + \frac{B'}{B\rho} \right) ds = \int_0^L K(s) ds$$
- And we can write our general expression for the chromaticity as
 
$$\xi = -\frac{1}{4\pi} \sum_i \beta_i \frac{1}{f_i} \Rightarrow \xi = -\frac{1}{4\pi} \oint \beta(s) K(s) ds$$

USPAS Fundamentals, June 4-15, 2018 E. Prebys, Accelerator Fundamentals: Off-Momentum Particles 22

## Chromaticity in Terms of Lattice Functions

- A *long* time ago, we derived the following constraint when solving our Hill's equation
 
$$w''(s) + K(s)w(s) - \frac{k}{w^3(s)} = 0 \Rightarrow (\sqrt{\beta})'' + K\sqrt{\beta} - \frac{1}{\beta^{3/2}} = 0$$

$$\begin{aligned} \beta(s) &= w^2(s) \\ \alpha(s) &= -\frac{1}{2}\beta'(s) \\ -\alpha^2 + \beta\gamma &= 1 \end{aligned}$$

$$(\sqrt{\beta})' = \frac{1}{2} \frac{1}{\sqrt{\beta}} \beta' = -\frac{\alpha}{\sqrt{\beta}}$$

$$(\sqrt{\beta})'' = -\frac{\alpha'}{\sqrt{\beta}} + \frac{1}{2} \frac{\alpha}{\beta^{3/2}} \beta' = -\frac{\alpha'}{\sqrt{\beta}} - \frac{\alpha^2}{\beta^{3/2}}$$

$$\Rightarrow K\beta^2 - \beta\alpha' - \alpha^2 = 1$$

Multiply by  $\beta^{3/2}$

- (We're going to use that in a few lectures), but for now, divide by  $\beta$  to get
 
$$K\beta = \frac{1+\alpha^2}{\beta} + \alpha' = \gamma + \alpha'$$
- So our general expression for chromaticity becomes
 
$$\xi = -\frac{1}{4\pi} \oint (\gamma(s) + \alpha'(s)) ds$$

USPAS Fundamentals, June 4-15, 2018

E. Prebys, Accelerator Fundamentals: Off-Momentum Particles

23

## Chromaticity and Sextupoles

- we can write the field of a sextupole magnet as
 
$$B(x) = \frac{1}{2} B'' x^2 \quad (\text{often expressed } b_2 x^2)$$
- If we put a sextupole in a dispersive region then off momentum particles will see a gradient
 
$$B'(x = D\delta) \approx B'' D \frac{\Delta p}{p_0}$$
 which is effectively like a position dependent quadrupole, with a focal length given by
 
$$\frac{1}{f_{eff}} = \frac{B''}{(B\rho)} LD \frac{\Delta p}{p_0}$$
- So we write down the tune-shift as
- Note, this is only valid when the motion due to momentum is large compared to the particle spread.

$$\Delta \nu = \frac{1}{4\pi} \beta \frac{1}{f_{eff}} = \frac{1}{4\pi} \frac{\beta B''}{(B\rho)} LD \frac{\Delta p}{p_0} \equiv \xi \frac{\Delta p}{p_0}$$

$$\Rightarrow \xi_S = \frac{1}{4\pi} \frac{\beta B''}{(B\rho)} LD$$